

Novel field theory phenomena from F theory and D-branes

J. Sonnenschein ^{a *}

^a School of Physics and Astronomy, Beverly and Raymond-Sackler Faculty of Exact Sciences, Tel-Aviv University, Ramat-Aviv, Tel-Aviv 69978, Israel

Talk presented in the 97 Karpacz winter school. We describe Sen's work on F theory on K3 and its reflection on the world-volume field theory on a D3-brane probe. Field theories on a multiple of probes are analyzed. We construct a 4d N=1 superconformal probe theory which is invariant under electric-magnetic duality via a compactification to six dimensions on $\mathbf{T}^6/\mathbf{Z}_2 \times \mathbf{Z}_2$.

1. Introduction

Recent developments in string theories, and in particular brane physics, have provided new magnifying glasses for observations on quantum field theories in various dimensions. In this series of talks presented in the Karpacz 97 winter school we have focused on exploring field theories on the world-volume of D-brane probes in F theory[1]. We have followed the analysis of A. Sen[2] on F theory on K3 and its reinterpretation in terms of a probe theory[4].

General background on D-branes, type II_B theory, orientifolds and F theory is presented in the introduction section. Section 2 is devoted to the analysis of F theory compactified on K3[2] and its relation to the field theory on a D3-brane probe[?]. A generalization of the latter to the theory on a multiple probe system that is put in various configurations with respect to the orientifold plane[5] is presented in section 3. Section 4 is devoted to the extraction of the properties of a probe theory in the neighbourhood of an intersecting orientifold planes of an F compactified on $\mathbf{T}^6/\mathbf{Z}_2 \times \mathbf{Z}_2$. The field content is found using a Gimon-Polchinski[3] type of analysis and a superpotential of this $N = 1$ theory, which admits a fixed line passing through the origin of the space of couplings, is written down. The last 2 sections are based on a work that was done in

collaboration with O. Aharony, S. Theisen and S. Yankielowicz[5],

1.1. A brief review of D-branes

A Dp-brane is a p dimensional extended object prescribed by the property that ends of open strings can lie on it, although open strings cannot exist in the bulk.[6] These open strings have Neuman boundary conditions (b.c) $\partial_n X^\mu = 0$ for $\mu = 0, 1, \dots, p$ and Dirichlet b.c $X^\mu = \text{constant}$ for $\mu = p + 1, \dots, 9$.

We summarize several properties of D-branes that we will make use of [7]: (i) The D-brane is a dynamical object. It can fluctuate in transverse directions. Its world-volume theory includes gauge fields and scalar fields. The latter correspond to the fluctuations. (ii) A Dp-brane couples to a A_{p+1} form. Thus type II_A includes even branes and II_B odd ones. (iii) A T-duality on S^1 interchanges the Dirichlet and Neuman b.c on the S^1 , so that a wrapped Dp-brane $\rightarrow$ D(p-1)-brane and unwrapped Dp-brane $\rightarrow$ D(p+1)-brane. (iv) a D-brane is a BPS state. It breaks half of the supersymmetries. (v) There is no force between infinite parallel D-branes at rest. (vi) N parallel D-branes can form a bound state at threshold.

1.2. A. brief review of type II_B string theory

We briefly summarize the bosonic fields and the symmetries of type II_B string theory.

1. Massless bosonic fields:

Neveu-Schwarz (NS-NS) sector- ϕ , $g_{\mu\nu}$, $B_{\mu\nu}^{(NS)}$.

*Work supported in part by the US-Israel Binational Science Foundation, by GIF – the German-Israeli Foundation for Scientific Research, and by the Israel Science Foundation.

Ramond-Ramond (RR) sector- a , $B_{\mu\nu}^{(RR)}$, $D_{\mu\nu\rho\sigma}^+$

2. Symmetries

I. Symmetries of the perturbative theory-

(i) $(-1)^{F_L}$

$$a \rightarrow -a; B_{\mu\nu}^{(RR)} \rightarrow -B_{\mu\nu}^{(RR)}; D_{\mu\nu\rho\sigma}^+ \rightarrow -D_{\mu\nu\rho\sigma}^+$$

(ii) Ω world sheet parity transformation-

$$a \rightarrow -a; B_{\mu\nu}^{(NS)} \rightarrow -B_{\mu\nu}^{(NS)}; D_{\mu\nu\rho\sigma}^+ \rightarrow -D_{\mu\nu\rho\sigma}^+$$

II. conjectured non-perturbative symmetry

$SL(2, \mathcal{Z})$ S-duality-

$$\lambda \rightarrow \frac{p\lambda + q}{r\lambda + s} \begin{pmatrix} B_{\mu\nu}^{(NS)} \\ B_{\mu\nu}^{(RR)} \end{pmatrix} \rightarrow \begin{pmatrix} p & q \\ r & s \end{pmatrix} \begin{pmatrix} B_{\mu\nu}^{(NS)} \\ B_{\mu\nu}^{(RR)} \end{pmatrix}$$

where

$$\lambda = a + ie^{-\phi}$$

$$p, q, r, s \in \mathcal{Z} \text{ with } ps - qr = 1.$$

1.3. A brief review of type II_B orientifolds

Type II_B defined on $\frac{\mathcal{M}}{G}$ where $\mathcal{M}$ -manifold and G - a group containing at least one element of the form

$$g_{s-t} \cdot g_{int} \cdot \Omega$$

where g_{s-t} is a space-time transformation, g_{int} - is an internal transformation and Ω is a world-sheet parity transformation.

Here are some examples of type II_B orientifolds

1. type II_B on $\frac{M^{9,1}}{\{1, \Omega\}}$ This is type I string theory on $M^{9,1}$.

2. type II_B on $M^{7,1} \times \frac{T^2}{\mathcal{I}_{89}(-1)^{F_L} \Omega}$ $\mathcal{I}_{89}$ transforms $x^8 \rightarrow -x^8$ $x^9 \rightarrow -x^9$

Upon performing T-duality on T^2 this is type I on $M^{7,1} \times T^2$.

3. type II_B on $M^{5,1} \times \frac{T^4}{(\mathcal{I}_{67}(-1)^{F_L} \Omega) \times \mathcal{I}_{89}(-1)^{F_L} \Omega}$

1.4. A brief review of F theory

In conventional compactifications of type II_B $\lambda = \text{constant}$.

F theory is defined in the following way. Let $\mathcal{M}$ be a manifold of dimension $d+2$ admitting an elliptic fibration. Let $\mathcal{B}$ be a (base) manifold of dimension d . $\mathcal{M}$ is obtained by erecting at every point of $\mathcal{B}$ a copy of a torus T^2 with its complex structure varying over $\mathcal{B}$.




Figure 1.

$$\text{F theory on } \mathcal{M} \equiv \text{type } II_B \text{ on } \mathcal{B}$$

with

$$\lambda = \tau(z) \neq \text{constant}$$

Note that when one moves along a closed cycle $\lambda = \tau(z)$ must come to its original value up to an $SL(2, \mathcal{Z})$ transformation. An example of elliptic fibration- elliptically fibered $K3$. In this case $\mathcal{M}$ - $K3$ and $\mathcal{B}$ - CP^1 . For z the coordinate of CP^1 the torus is parametrized by the complex coordinates x and y

$$y^2 = x^3 + f_8(z)x + g_{12}(z) \quad (1)$$

$f_8(z)$ and $g_{12}(z)$ are polynomials of degree 8 and 12 respectively.

The J invariant of T^2 is given by

$$j(\tau(z)) = \frac{4(24f)^3}{27g^2 + 4f^3} \quad (2)$$

F theory on $K3$ is equivalent to type II_B on CP^1 with $\lambda = \tau(z)$. We enlist here several **Dualities of F theory** that will be useful in the discussions below.

1. F theory on $\frac{T^2}{\mathbb{Z}_2} \Leftrightarrow$ type I (heterotic) $SO(32)$
2. F theory on $\mathcal{M} \times S_1 \Leftrightarrow$ type II_B on $\mathcal{B} \times S_1$ with $\lambda(z) = \tau(z)$ of the fibers. Recall that type II_B on S_1 with radius R is dual to type II_A on S_1 with radius $\frac{1}{R}$. Furthermore, type II_A on S_1 is equivalent to M-theory on T^2 , so that λ of the type II_B theory is identified with τ of the latter T^2 . We thus conclude that

F theory on $\mathcal{M} \times S_1 \Leftrightarrow$ M-theory on $\mathcal{M}$

3. From (1) F theory on $K3 \times S_1 \Leftrightarrow$ M-theory on $K3 \Leftrightarrow$ Heterotic on T^3 .

In a similar manner, F theory on $K3 \Leftrightarrow$ Heterotic on T^2 .

Since the heterotic theory on $T^2 \Leftrightarrow$ type I on $T^2 \Leftrightarrow$ type II_B on $\frac{T^2}{\mathbb{Z}_{89}}$ one finds

F theory on $K3 \Leftrightarrow$ type II_B on $\frac{T^2}{\mathbb{Z}_{89}}$

4. F theory on $K3$ fibered CY $\Leftrightarrow$ heterotic on $K3$.

2. Sen's model- F-theory on K3- space-time and worldvolume theories

Elliptic fibration of $K3$ is described by eqn. (1). The idea of Sen was to look for a point in the F-theory moduli space where

$\tau(z)$ is independent of z and $Im(\tau)$ is large

This should imply a conventional perturbative string compactification, and as will be shown below a conformal invariant field theory on a “probe” worldvolume.

A constant $\tau(z)$ means also

$$j(\tau(z)) = \text{constant} \rightarrow \frac{f^3}{g^2} = \text{constant} \quad (3)$$

This condition is obeyed if one takes

$$f = \alpha \phi(z)^2, \quad g = \phi(z)^3 \quad (4)$$

In that case $j(\tau(z)) = \frac{4(24\alpha)^3}{27+4\alpha^3} = \text{constant}$ where α can be adjusted so that $Im(\tau)$ is large.

Singularities of $j(\tau)$ at

$$\Delta = (27g^2 + 4f^3) = 0 \quad (5)$$

For the ansatz (4) and with $\phi \sim \prod_{i=1}^4 (z - z_i)$

$$\Delta \sim \prod_{i=1}^4 (z - z_i)^6 \quad (6)$$

Metric on the base CP^1

In general the metric is given by [10]

$$ds^2 = F(\tau, \bar{\tau}) \prod_{i=1}^{24} (z - z_i)^{-\frac{1}{12}} \prod_{i=1}^{24} (\bar{z} - \bar{z}_i)^{-\frac{1}{12}}$$

So for the point in moduli space we consider it takes the form

$$ds^2 = F(\tau, \bar{\tau}) \prod_{i=1}^4 (z - z_i)^{-\frac{1}{2}} \prod_{i=1}^4 (\bar{z} - \bar{z}_i)^{-\frac{1}{2}}$$

This can be brought to the form of a **Flat metric**. Define

$$dw = \prod_{i=1}^4 (z - z_i)^{-\frac{1}{2}} dz$$

then

$$ds^2 = dw d\bar{w}.$$

Near $z = z_i$, $w \sim (z - z_i)^{-\frac{1}{2}}$, namely, there is a **Conical deficit angle of π** at each of the four z_i .

Naively, we may conclude that

$$\mathcal{B} = \frac{T^2}{\mathbb{Z}_{89}} \quad (7)$$

But there is a subtlety, lets check again

$$y^2 = x^3 + \alpha x \prod_{i=1}^4 (z - z_i)^2 + \prod_{i=1}^4 (z - z_i)^3 \quad (8)$$

going around an orbifold point like $z = z_i$

$$\begin{aligned} (z - z_i) &\rightarrow e^{2\pi i} (z - z_i) \\ x &\rightarrow e^{2\pi i} x = x \\ y &\rightarrow e^{3\pi i} y = -y \end{aligned}$$

which is

$$\text{An } \text{SL}(2, \mathbb{Z}) \text{ trans. } \begin{pmatrix} -1 & \\ & -1 \end{pmatrix} \text{ on the } T^2 \text{ fiber}$$

The transformation $\begin{pmatrix} -1 & \\ & -1 \end{pmatrix}$ corresponds to

$$(-1)^{F_L} \Omega$$

Thus, the the base is infact an **Orientifold**

$$\mathcal{B} = \frac{T^2}{(-1)^{F_L} \mathcal{I}_{89} \Omega}.$$

Under this projection $z \rightarrow -z$, so define $u = z^2$ which is single valued on the orientifold.

The orientifold

(1) preserves half of the space-time supersymmetries of type II_B .

(2) It carries a RR topological charge -4 [6]




Figure 2.

$$\oint_C da = -4 \quad (9)$$

where C is a contour around $u = 0$

D7-brane

(1) preserves half of the space-time supersymmetries of type II_B .

(2) It carries a RR topological charge $+1$

To cancell the RR charges of the 4 orientifolds the theory must have **16 D7-branes** see fig. 4.

Since we have a configuration with




Figure 3.




Figure 4.

$\lambda = \text{constant} \rightarrow a = \text{constant}$ the RR charge must be cancelled **locally** on the orientifold (fig. 5)

There is an **Enhanced symmetry at singularities** that can be viewed in F-theory picture and in the orientifold picture.

Start with F-theory picture

$$y^2 = x^3 + \alpha x \prod_{i=1}^4 (z - z_i)^2 + \prod_{i=1}^4 (z - z_i)^3 \quad (10)$$

Near $z = z_1$

$$\tilde{y}^2 = \tilde{x}^3 + \alpha \tilde{x} (z - z_1)^2 + (z - z_1)^3 \quad (11)$$

which means that there is a

$$D_4 \text{ type singularity} \rightarrow SO(8)^4 \text{ gauge symmetry}$$

Now the orientifold description Let us construct the string configuration in several steps: (i) From strings starting and ending on the 7Dbrane fig. 6. one finds




Figure 5.

$N = 1$ 8D wv theory with $U(1)$ gauge field and a complex scalar that corresponds to the location of the D-brane in the z plane.

(ii) When two D-branes collapse one gets (fig. 7)




Figure 6.

$N = 1$ 8D wv theory with $U(2)$ gauge fields and a complex scalar in the adjoint.

(iii) When all the four z_i are equal (fig. 8) then the

symmetry is enhanced to $U(4)$

(iv) when all the four $z_i = 0$ (fig. 9)

there are additional strings (and hence massless vectors) between the 7branes at z_i and their mirrors at $-z_i$. Now the symmetry is enhanced to $U(8)$

(v) But the generator of $Z_2 = (-1)^{F_L} \mathcal{T}_{89} \Omega$ acts on the generators T^a of the $U(8)$.

Only those that commute with it survive. Un-




Figure 7.




Figure 8.

der Z_2

$$T^a \rightarrow_{Z_2} - \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix} (T^a)^T \begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix}$$

the various factors of this transformations are (i)-sign: change the sign of the oscillator A_{-1}^μ that creates open strings from the vacuum.

(ii) Transpose: effect of orientation reversal

(iii) $\begin{pmatrix} 0 & I_4 \\ I_4 & 0 \end{pmatrix}$: effect of exchange of Dbranes and their images. This breaks

$$U(8) \rightarrow SO(8)$$

Let us now move away from the special point in the moduli space by switching background fields in both the orientifold and F-theory descriptions.

Deforming away from the special point-

Orientifold description- At each of the orientifolds the 7D-branes can be moved away from the orientifold. Near one particular orientifold the

theory is type II_B on $\frac{\mathcal{R}^2}{(-1)^{F_L} \mathcal{I}_{89}\Omega}$ and the picture is like that of the neighborhood of a D7-brane in fig. 5.




Figure 9.

The background field λ

$$\lambda(z) = \lambda_0 + \frac{1}{2\pi i} \left(\sum_{i=1}^4 \ln(z - z_i) - 4 \ln z \right)$$

It is clear that **close to** $z = 0$ **solution not consistent** since $Im(\lambda) < 0$.

F-theory description

Around the orbifold point

$$y^2 = x^3 + f(z)x + g(z)$$

where $f(z)$ and $g(z)$ are polynomial of degree 2 and degree 3 respectively.

The zeros of Δ are now at 6 singular points




Figure 10.

The total number of parameters is

$$\begin{aligned} & 3 \text{ from } (f) + 4 \text{ from } (g) \\ & -1 \text{ from rescaling of } x, y - 1 \text{ shift of } z = 5 \end{aligned}$$

Recall the Seiberg-Witten solution for $N = 2$ $SU(2)$ gauge symmetry with $N_f = 4$ is parametrized by

$$\tau(z, m_1, m_2, m_3, m_4, \tau_0)$$

where $z = \langle \phi^2 \rangle$, m_i are quark masses and τ_0 is the classical coupling. It is now natural to use the SW parameters z, τ_0, m_i to label the F-theory background.

In this description the singularities are associated with massless quarks, massless monopole and massless dyon.

	orientifold	F-theory
parameters	$\lambda_0, z_1, z_2, z_3, z_4$	$\tau_0, m_1, m_2, m_3, m_4$
back-ground λ	$\lambda(z) = \lambda_0 + \frac{1}{2\pi i} \left(\sum_{i=1}^4 \ln(z - z_i) - 4 \ln z \right)$	$\lambda(z) = \tau(\tau_0, m_i)$
asymptotic λ	λ_0	τ_0
BPS spectrum	z_i	m_i^2

The orientifold solution for $\lambda(z)$ is the same as the SW solution for large z (u)

$$\tau(z) = \tau_0 + \frac{1}{2\pi i} \left(\sum_{i=1}^4 \ln(z - m_i^2) - 4 \ln z \right)$$

The singularities $\ln(z - m_i^2)$ are due to the massless quarks and the $4 \ln z$ term is associated with massless $SU(2)$ gauge bosons which splits into two singularities in the full solution. The comparison between the orientifold and the F theory picture is summarized in the table above.

We can now conclude that

F-theory provides the quantum corrected version of the orientifold background in the form of **Seiberg-Witten** solution

Question:

Is it a coincidence that the SW theory emerged?

2.1. SW solution as the wv theory on a 3B probe

Let us study the effective Lagrangian on a probe that moves in the background of the F-theory on K3.

Question: What probe should one use? 3Brane is the most natural probe since: (1) 3Brane is invariant under $SL(2, \mathcal{Z})$, so the wv theory should also be invariant under $SL(2, \mathcal{Z})$ duality, (2) 7Brane parallel to a 3Brane probe breaks 1/2 of the supersymmetries so that $\rightarrow N = 2$ in 4D.

Another interpretation of the 3Brane probe is as a **wrapped 5Brane**. Recall F-theory on K3 is dual to type I on T^2 . Take a 5Brane of type I. Perform a T-duality on the T^2 which interchanges $N \Leftrightarrow D$ boundary conditions, and you find a 3D-Brane of type I'. The Chan-Paton $SU(2)$ gauge group of the 5Brane of type I is inherited by the 3Brane of the type I'. The comparison between the two dual pictures is summarized in the following table[4].

type I 5Brane	type I' 3Brane
Wilson lines around cycles of T^2 $SU(2) \rightarrow U(1)$	3Brane position on $\frac{T^2}{\mathbb{Z}_2}$ away from orientifold $SU(2) \rightarrow U(1)$
position of 7Brane $R'_8 \alpha_{i,8} + i R'_9 \alpha_{i,9}$ position of 3Brane $R'_8 A_8 + i R'_9 A_9 = \begin{pmatrix} w & 0 \\ 0 & -w \end{pmatrix}$	position of 7Brane m_i position of 3Brane $z = w^2$

where the notation in this table is that of ref. [4]. The strings stretched from the 3Brane to the 7Brane induce fundamental matter hypers with mass $m_i \pm w$.

Let us summarize the correspondence between

the F theory and the probe theory:

(i) The set-up on the 3Brane wv is that of $SU(2)$ SW theory with four hyper-multiplets in the fundamental rep.

(ii) The F-theory $\lambda \rightarrow$ probe gauge coupling τ .

(iii) The fact that $\lambda = \text{constant} \rightarrow$ into conformal invariance of the probe theory.

(iv) type II_B $SL(2, \mathcal{Z})$ invariance $\rightarrow$ $SL(2, \mathcal{Z})$ invariance of $SU(2)$ with $N_f = 4$

3. The world-volume field theory on a multiple of 3-branes probes in 8 dimensions

Consider again F theory on K3. Recall that F theory on K3 is dual to type I on T^2 . Now wrap N_c 5B around the T^2 . Let us analyze the properties of a world-volume theory on the multiple probes [14].

(i) *Supersymmetry*- 8 susy charges $N = 1$ in 5+1 dim $\rightarrow N = 2$ in 3B 3+1 wv.

(ii) *Gauge symmetry*- $USP(2N_c)$ [8]

Hypermultiplets- (a) A - antisymmetric rep. $N_c(2N_c - 1)$ which is irreducible $N_c(2N_c - 1) - 1 \oplus$ singlet.

(b) $N_f = 4$ Q^i , “quarks” in the fundamental rep. $2N_c$ which emerge in F theory from strings between 3B and the four 7B at the orientifold point.

(iii) *Global symmetry*- $SO(8)$ global symmetry (local symmetry in space-time)

(iv) *Superpotential*- $W = Q^i X Q^i + A X A + m_{ij} Q^i Q^j$

(v) *Mass terms*- m_{ij} corresponds to the complex scalar 28 antisymmetric rep. of the space-time $SO(8)$ local sym. No other scalars in space-time $\rightarrow$ No mass term to A .

(vi) β function $(2N_c + 2) - (2N_c - 2) - 4 = 0$

(vii) *Expectation values*- (a) scalars of the antisymmetric rep which correspond to the motion in x_4, x_5, x_6, x_7 . (b) scalars of vector multiplet associated with Wilson loops of the gauge fields around T^2 .

(viii) *Moduli space of Coulomb branch*-

$$\frac{(SU(2)^k)}{S_k}$$

permutation symmetry S_k part of the $USP(2N_c)$ Weyl group.

3.1. Flows of the $USP(2N_c)$ theory

$$(1) USP(2N_c) \rightarrow U(1)^{N_c}$$

$$\langle X \rangle = \text{diag}(a_1, a_2, \dots, a_k, -a_1, -a_2, \dots, -a_k)$$

where a_i^2 corresponds in the type II_B picture to the location of the i^{th} 3B on $\frac{T^2}{Z_2}$ from the orientifold.




Figure 11.

$$(2) USP(2N_c) \rightarrow SU(2)^{N_c}$$

$$\langle A \rangle = \begin{pmatrix} 0 & 0 & \cdots & 0 & b_1 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 & 0 & b_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 0 & 0 & \cdots & b_k \\ -b_1 & 0 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ 0 & -b_2 & \cdots & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & -b_k & 0 & 0 & \cdots & 0 \end{pmatrix}.$$

b_i^2 corresponds in the type II_B picture to the location of the i^{th} 3B in x_4, x_5, x_6 and x_7 .

In the limit of all b_i being very large only singlet remanants from A survives so it flows to N_c copies of Seiberg-Witten theory of $SU(2)$ with $N_f = 4$.

$$(3) USP(2N_c) \rightarrow U(N_c); N = 4$$

The N_c 3B are together but very far from the orientifold point [9].

One can deduce from the probe theory interesting properties of the SW theories [5] To summarize:

The Coulomb phase of wv field theory of multiple p -branes is just N_c copies (up to global identifications) of the Coulomb phase for a single p -brane.



Figure 12.

4. Construction of $N = 1$ superconformal 4D probe theory with electric-magnetic duality symmetry

Consider as in Sen's model[2] compactifications of F- theory on CY manifolds which have a constant (weak coupling) value of τ .

The simplest case is the elliptic fiber over a base $\mathbf{CP}^1 \times \mathbf{CP}^1$ [11]. In analogy to (1) the toric fibers




Figure 13.

are described by

$$y^2 = x^3 + f(z_1, z_2)x + g(z_1, z_2),$$

where z_1 and z_2 label the two $\mathbf{CP}^1$ s of the base.

$$\Delta = 4f^3 + 27g^2 = 0$$

At a degeneration of the fibers, which is a location of type IIB 7-brane in the compact dimensions, τ has a non-trivial monodromy when going around it.

(i) in F-theory

$$\tau(z_1, z_2) = \text{constant} \rightarrow \frac{f^3}{g^2} = \text{constant}$$

A solution (though not the most general solution) which generalizes (4) is

$$f(z_1, z_2) = \alpha \phi_1(z_1)^2 \phi_2(z_2)^2; \quad g(z_1, z_2) = \phi_1(z_1)^3 \phi_2(z_2)^3$$

where ϕ_1 and ϕ_2 are general polynomials of degree four

$$\phi_1(z_1) = \prod_{i=1}^4 (z_1 - z_1^{(i)}); \quad \phi_2(z_2) = \prod_{i=1}^4 (z_2 - z_2^{(i)}).$$

At this point in moduli space $\tau = \text{constant}(\alpha)$ and there are D_4 singularities at $z_1 = z_1^{(i)}$ and at $z_2 = z_2^{(i)}$.

4.1. Space-time theory at the intersection of the two D_4 singularities

In general the spacetime field theory at an intersection of two D_4 singularities is not well understood[12,15]. Let us try to interpret this point in moduli space as an orientifold of the type IIB theory. Around each of the points $z_1 = z_1^{(i)}$ (for constant z_2) there is an $SL(2, \mathbf{Z})$ monodromy of the form

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Locally, this monodromy can be interpreted as an orientifold of the type IIB theory by $(-1)^{F_L} \cdot \Omega$, accompanied by four 7-branes to cancel the RR charge.

Each point $z_1 = z_1^{(i)}$ on the first $\mathbf{CP}^1$ factor carries a deficit angle of π , all four of them together deforming the two $\mathbf{CP}^1$ to $\mathbf{T}^2/\mathbf{Z}_2$. Thus, the theory looks like

$$\text{type IIB on } \frac{\mathbf{T}^4}{\mathbf{Z}_2 \times \mathbf{Z}_2'}$$

where

$$z_2 = \mathcal{I}_{67} \Omega(-1)^{F_L}; \quad z_2' = \mathcal{I}_{89} \Omega(-1)^{F_L}.$$

Naively, this corresponds to the orientifold of F-theory on $\frac{\mathbf{T}^6}{\mathbf{Z}_2 \times \mathbf{Z}_2'}$.

However, this orientifold compactification **cannot be identified directly with the F-theory compactification**[?]

Generally, in F-theory compactification

$$B_{\mu\nu}^{NS} = B_{\mu\nu}^{RR} = 0$$

our case there is **discrete non-vanishing $B_{\mu\nu}$ fields** (related to *discrete torsion*) *How do we know?* At the intersection of two D_4 singularities the intersection point can be blown up to get an additional 2-cycle. In the naive F-theory compactification 3-brane can be wrapped over the vanishing 2-cycle, which should mean that there are $\Rightarrow$ **tensionless strings** living on the intersection of the D_4 singularities. Our orientifold construction, on the other hand, involves a well-defined weakly coupled conformal field theory, whose low-energy spectrum is well-defined and can not include such tensionless strings. Discrete $\mathbf{Z}_2$ symmetries force the value of both 2-form tensor fields (integrated over the vanishing 2-cycle) to be either 0 or 1/2 (modulo 1).

A non-zero value for either (or both) of these fields breaks the $SL(2, \mathbf{Z})$ symmetry to a discrete $\Gamma(2)$ subgroup.

$$\Gamma(2); \{s = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, t^2 = \begin{pmatrix} 1 & 2 \\ -2 & 1 \end{pmatrix}\}$$

There are two possible models:

- (i) Integrated $B^{NS} = \frac{1}{2}, B^{RR} = 0 \rightarrow$ **Sen's $SU(4)$ model** [13]
- (ii) Integrated $B^{NS} = B^{RR} = \frac{1}{2} \rightarrow$ **$SO(8)$ model**[5]

4.2. The space-time theory of the $SO(8)$ model

The superfield content which one deduce from the various string sectors are given by:

- (1) *Untwisted closed string sectors* :
supergravity multiplet + one tensor multiplet + 4 hypermultiplets.

Open string sector:

- (i) 7-7 strings $\rightarrow SO(8)^4$ vector multiplets.
- (ii) 7'-7' strings $\rightarrow SO(8)^4$ vector multiplets.
- (iii) 7-7' no massless states.

- (2) *Twisted closed string sectors* :

A tensor multiplet at each intersection of fixed points (i.e. at each fixed point of both $\mathbf{Z}_2$'s), accounting for a total of **16 additional tensor multiplets**. At each intersection of D_4 singularities there is a possibility of blowing up a point to

get an additional 2-cycle After all these blow-ups we get

F-theory on a smooth ($h_{11} = 51, h_{21} = 3$) **Calabi-Yau manifold.**

which includes [16–18]:

$$n_T = h_{11}^B - 1 = 18 - 1$$

$$n_{H^0} = h_{21}^M + 1 = 3 + 1 = 4$$

$$r(V) = h_{11}^M - h_{11}^B - 1 = 51 - 18 - 1 = 32$$

This set of 6D $N = 1$ multiplets obey the anomaly cancellation condition

$$29n_T + n_H - n_V = 293$$

Note again, that the spacetime theory we found (using the orientifold construction) is not the same as the theory we supposedly started with, which was F-theory compactified on an elliptic fibration over $\mathbf{CP}^1 \times \mathbf{CP}^1$.

4.3. The space-time theory of the $SU(4)$ model

[13]

Untwisted closed string sectors :

supergravity multiplet + one tensor multiplet + 4 hypermultiplets.

Open string sector:

(i) 7-7 strings $\rightarrow U(4)^4$ vector multiplets + 2×6 hypers for each vector.

(ii) 7'-7' strings $\rightarrow U(4)^4$ vector multiplets + 2×6 hypers for each vector.

(iii) 7-7' (4, 4) hyper multiplets.

Twisted closed string sectors :

16 hypers

$$n_T = 1, \quad n_H = 372 \quad n_V = 128$$

4.4. 3B probe wv field theory at the intersection

The probe theory is expected (i) to have only $N = 1$ **supersymmetry**. (since the 7-branes intersect transversely at this point and each breaks a different half of the wv $N = 4$ supersymmetry,) (ii) to be invariant under conformal and ($\Gamma(2)$) electric-magnetic duality symmetry, and (iii) to have two deformations, corresponding to turning on z_1 or z_2 , which should cause it to flow to the $N = 2$ $SU(2)$ gauge theory with 4 quark hypermultiplets.

4.5. Fields from the 3-brane strings

Let us begin by computing the fields corresponding to strings which stretch out from a 3-brane and fold back to the same brane.

The 3-brane has 3 images under $\mathbf{Z}_2 \times \mathbf{Z}_2$, so that every open string state is enhanced to a 4×4 matrix (as in Gimon-Polchinski[3] (fig. 14))

imposing the orientifold restrictions.




Figure 14.

The γ matrices[3] corresponding to each of the generators in the $\mathbf{Z}_2 \times \mathbf{Z}_2$ may be chosen to be

$$\gamma_{\Omega_1} = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{pmatrix} \quad \gamma_{\Omega_2} = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix}$$

and the orbifold matrix is then necessarily

$$\gamma_T = \gamma_{\Omega_1} \gamma_{\Omega_2} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}. \quad (12)$$

These matrices are hermitian and unitary, and they correctly reflect the two $\mathbf{Z}_2$ actions on the compact coordinates (z_1, z_2) .

The wave-function matrices of states, M_{ij} , must then satisfy conditions of the type

$$\begin{aligned} M &= \pm \gamma_{\Omega_1} M^T \gamma_{\Omega_1}^{-1} \\ M &= \pm \gamma_{\Omega_2} M^T \gamma_{\Omega_2}^{-1} \\ M &= \pm \gamma_T M \gamma_T^{-1} \end{aligned} \quad (13)$$

where the signs are determined by the transformation properties of the relevant state.

For the gauge fields- $-,-,+,$

For the chiral superfield $X_{6,7} - -, +, -,$

For $X_{8,9} - +, -, -$

For $X_{4,5} - +, +, +.$

Note that in flat space the 3-brane field theory involves an $N = 4$ vector multiplet, containing an $N = 1$ vector multiplet and three $N = 1$ chiral multiplets corresponding to the coordinates $X_{4,5,6,7,8,9}$ of the 3-brane. In the presence of the orientifold, each of these fields is enhanced into a 4×4 matrix with different constraints.

(i) Vector superfields

The relations imposed by (13) on the components M_{ij} of gauge fields are

$$\begin{aligned} M_{11} &= -M_{22} = -M_{33} = M_{44} \\ M_{14} &= M_{41} = M_{23} = M_{32} \\ M_{12} &= -M_{43} \quad M_{13} = -M_{42} \\ M_{21} &= -M_{34} \quad M_{31} = -M_{24}. \end{aligned} \quad (14)$$

A basis of 6 matrices that obey these relations is

$$\begin{aligned} Z_1 &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} & Z_2 &= \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix} \\ W_1 &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{pmatrix} & W_2 &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix} \\ W_3 &= \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix} & W_4 &= \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{pmatrix} \end{aligned}$$

When we take z_1 to infinity all 1-2,1-4,2-3 and 3-4 strings become infinitely massive, (fig. 15) so we can drop those wave-function matrices with non-zero entries in those positions. This leaves Z_1, W_2 and W_3 , which generate an $SU(2)$ **sub-algebra**.

It is now straightforward to check that the matrices

$$\begin{aligned} W_2^+ &= \frac{1}{4}(W_1 + W_2); \quad W_1^- = \frac{1}{4}(W_1 - W_2); \\ W_1^+ &= \frac{1}{4}(W_3 + W_4); \quad W_2^- = \frac{1}{4}(W_3 - W_4) \\ Z^\pm &= \frac{1}{4}(Z_1 \pm Z_2) \end{aligned}$$

obey the following commutation relations

$$[Z^+, W_2^+] = -iW_1^+; \quad [Z^+, W_1^+] = iW_2^+;$$

$$\begin{aligned} [W_2^+, W_1^+] &= -iZ^+ \\ [Z^-, W_1^-] &= iW_2^-; \\ [Z^-, W_2^-] &= -iW_1^-; \quad [W_1^-, W_2^-] = iZ^- \\ [Z^+, W_1^-] &= 0; \quad [Z^+, W_2^-] = 0; \\ [Z^-, W_2^+] &= 0; \quad [Z^-, W_1^+] = 0; \quad [Z^+, Z^-] = 0 \\ [W_2^+, W_1^-] &= 0; \quad [W_2^+, W_2^-] = 0; \\ [W_1^+, W_1^-] &= 0; \quad [W_1^+, W_2^-] = 0. \end{aligned}$$

Thus, we see that the gauge fields span an $SU(2) \times SU(2)$ **algebra**.




Figure 15.

(ii) X_{67} chiral multiplet

Next consider the implications of [?] on the scalar fields X_{67} . The constraints on the matrix components are now

$$\begin{aligned} M_{11} &= -M_{22} = M_{33} = -M_{44} \\ M_{14} &= -M_{41} = -M_{23} = M_{32} \\ M_{12} &= +M_{43} \quad M_{13} = M_{42} = 0 \\ M_{21} &= +M_{34} \quad M_{31} = M_{24} = 0 \end{aligned}$$

A basis of hermitian matrices that obey the rela-

tions is the following

$$A_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix} \quad A_2 = \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}$$

$$A_3 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \quad A_4 = \begin{pmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \end{pmatrix}.$$

Note that now if we take z_1 to infinity (naively) we are left only with A_1 , while if we take z_2 to infinity (naively) we remain with A_1, A_3 and A_4 which are in the adjoint representation of the remaining $SU(2)$, and thus we go over to the picture of Sen, as expected.

Define now the matrices

A_{-+}, A_{++}, A_{--} and A_{+-}

by the equations

$$A_{++} = A_3 - iA_4; \quad A_{-+} = A_1 + iA_2; \quad A_{+-} = A_1 - iA_2 \text{ and } A_{--} = A_3 + iA_4.$$

It can easily be checked that

$A = (A_{++}, A_{-+}, A_{+-}, A_{--})$ are in the

$(2, 2)$ **representation of $SU(2) \times SU(2)$** For instance, we have

$$[Z^+, A_{++}] = \frac{1}{2}A_{++} \quad [Z^+, A_{-+}] = -\frac{1}{2}A_{-+}$$

$$[W_2^+, A_{++}] = -\frac{1}{2}A_{-+} \quad [W_2^+, A_{-+}] = -\frac{1}{2}A_{++}$$

$$[W_1^+, A_{++}] = \frac{i}{2}A_{-+} \quad [W_1^+, A_{-+}] = -\frac{i}{2}A_{++}$$

and so on.

(iii) X_{89} **chiral multiplet**

For the relations on the X_{89} chiral superfield we would find similar (but not identical) results to those of X_{67} (with the second and third rows and columns of all matrices interchanged.)

Thus, this chiral superfield, that we denote by

$$B = (B_{++}, B_{-+}, B_{+-}, B_{--})$$

, is also in the $(2, 2)$ **representation of $SU(2) \times SU(2)$**

As a consistency check, we verify that a VEV for X_{67} , for instance, indeed breaks the gauge

symmetry to a diagonal $SU(2)$. For instance, taking A_1 to have a VEV would leave exactly the matrices Z_1, W_2 and W_3 (given above), which commute with it, as expected. The A 's would also all become massive except A_1 (since they do not commute with it), again as expected (since after moving along the flat direction we should have just a single scalar).

(iv) X_{45} **chiral multiplet**

The relations for X_{45} are

$$M_{11} = M_{22} = M_{33} = M_{44}$$

$$M_{14} = M_{41} = -M_{23} = -M_{32},$$

and their solutions are spanned by the two singlets S_1 and S_2

$$S_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad S_2 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$

4.6. Fields form the strings between the 3-brane and 7-branes

The γ_Ω matrices for the 7-branes are all proportional to the identity matrix. The constraint on the 7-brane gauge fields is then just

$$M = -M^T,$$

giving an anti-symmetric matrix corresponding to an $SO(8)$ space-time gauge symmetry, since there are 8 7-branes when including the Z_2 partners.

Next, we should use these matrices to analyze the wave function of the 3-7 strings (as in [3]). The 7-brane side is trivial, so the orbifold/orientifold projections just mix the various 3-branes according to the γ_Ω matrices. For instance taking the $3_1 - 7$ state (where 3_1 is the first 3-brane) via γ_{Ω_1} to i times the $7 - 3_2$ state (with opposite orientation!), via γ_{Ω_2} to i times the $7 - 3_3$ state, and via γ_T to minus the $3_4 - 7$ state. A basis for the states going to a specific 7-brane can thus be chosen to be

$$D_u^+ = (1, 0, 0, -1); \quad D_d^+ = (0, 1, 1, 0).$$

Using the matrices we found above for the gauge fields, it is easy to check that these are doublets under $SU(2)_+$, and singlets with respect

to $SU(2)_-$. The corresponding chiral superfields are thus in $(2, 1)$ **representation of $SU(2)_+ \times SU(2)_-$** . For the other group of 7-branes we can then do the same thing, but with minus the identity matrix for some of the relevant 7-7 γ_Ω matrices (say for γ_{Ω_2} and γ_T). Then, the basis comes out to be

$$D_u^- = (1, 0, 0, 1); \quad D_d^-(0, 1, -1, 0).$$

which is in the $(1, 2)$ **representation of $SU(2)_+ \times SU(2)_-$** .

The summary of the field content is

A_μ — $SU(2) \times SU(2)$ **gauge fields**.

$X_{8,9} \equiv B$ — **chiral field in $(2, 2)$ $A^2 = z_2$**

$X_{6,7} \equiv A$ — **chiral field in $(2, 2)$ $A^2 = z_1$**

$X_{4,5} \equiv (S_1, S_2)$ — **singlet chiral fields**

Q^i ($i = 1, \dots, 8$)— **chiral quarks in $(2, 1)$**

q^i ($i = 1, \dots, 8$)— **chiral quarks in $(1, 2)$**

Note about this particle spectrum.

(i) It is manifestly not $N = 2$ supersymmetric (since there is no chiral multiplet in the adjoint representation)

(ii) $\langle A \rangle \neq 0$ ($\langle B \rangle \neq 0$) breaks $SU(2) \times SU(2) \rightarrow SU(2)$ -

Three components of A (B) are swallowed by the Higgs mechanism, and we remain with an adjoint chiral multiplet (coming from B (A)) and additional singlets, as expected.

(iii) A natural interpretation of S_1 S_2 is that at the intersection point the 3-brane can split into two half-3-branes which can move independently.

(iv) **Vanishing one loop β function**

(since there are 12 ($N_f = 6$) doublets of each $SU(2)$).

4.7. The Superpotential

Let start with some general points:

(i) In principle, it should be possible to compute this superpotential from the string theory analysis, but we have not performed this computation.

(ii) Since we have only $N = 1$ supersymmetry, the superpotential in general receives non-perturbative quantum corrections.

(iii) The superpotential has to obey the following main constraints:

1) The $(SO(8))$ theory should have (at least) an $SO(8) \times SO(8)$ global symmetry. The $(SU(4))$

theory should have (at least) an $SU(4) \times SU(4)$ global symmetry.

2) The theory should flow to the $N = 2$, $SU(2)$, $N_f = 4$ theory upon giving a VEV to A or to B .

3) The theory should have a fixed line passing through weak coupling [19,20].

4) The quarks q^i (Q^i) have to become massive when we give a VEV to A (B)

Construction of the superpotential:

(i) **Superpotential for the quarks** The vanishing of the β function associated with the gauge coupling β_g and with the superpotential couplings β_{h_n} requires that

$$A_g = 3C_2(G) - \sum_i T(r_i) + \sum_i T(r_i)\gamma_i = 0 \quad (15)$$

$$A_{h_n} = (n-3) + \frac{1}{2} \sum_k \gamma_k = 0 \quad (16)$$

$$(17)$$

where in the first equation the sum is over all the fields in the theory, and in the second only over the fields in the particular term of the superpotential.

A fixed line passing through weak coupling demands that these equations are not all independent [19]. It is easy to check that $n = 3$ and $\gamma_k = 0$ is a solution for this condition. In fact in [20] it was shown from string arguments that indeed all the anomalous dimensions vanish.

Combining this solution with the requirements 1), 2) and 4) one ends up with the following superpotentials

For the $SU(4)$ model [13]

$$W = QA\tilde{Q} + qB\tilde{q} \quad (18)$$

and for the case of $SO(8)$ [5]

$$W = \frac{QABQ}{\sqrt{A^2}} + \frac{qABq}{\sqrt{B^2}} \quad (19)$$

In spite of the fact that in both cases these superpotentials obey all the requirements, it is clear that only for the $SU(4)$ model it is an adequate solution. In the second case one encounters singularities in the limits of $A \rightarrow 0$ and $B \rightarrow 0$ which cannot be smoothed.

(ii) **Construction of the superpotential for S_1, S_2**

(a) S_1 - the location of the 3-brane in $X_{4,5}$ coordinates should be decoupled.

(b) S_2 (corresponding to splitting the 3-brane at the orientifold point) should be massless only when A and B are both zero.

(c) A and B become massive once S_2 is turned on, as expected, since when the 3-brane has split we cannot move it away from the orientifold point.

(d) In the “Coulomb branch” $\langle A^2 \rangle$ and $\langle B^2 \rangle$ should correspond to flat directions but not AB .

(e) $\langle A \rangle$ -three of its components are swallowed by the Higgs mechanism, and the other remains massless and parametrizes the flat direction corresponding to the flow (it is the $N = 2$ partner of S_1). Three of the components of B remain massless and become an adjoint field X of the remaining $SU(2)$, but the remaining component, as well as S_2 , should become massive (since we have no corresponding fields in the $N = 2$ theory) The final result for this part of the action is

$$W = S_2 AB$$

Acknowledgements We would like to thank A. Brandhuber, V. Kaplunovsky and especially A. Sen for useful conversations.

REFERENCES

1. C. Vafa, “Evidence for F-Theory”, Nucl. Phys. **B469** (1996) 403, hep-th/9602022
2. A. Sen, “F Theory and Orientifolds”, Nucl. Phys. **B475** (1996) 562, hep-th/9605150
3. E. G. Gimon and J. Polchinski, “Consistency Conditions for Orientifolds and D-Manifolds”, Phys. Rev. **D54** (1996) 1667, hep-th/9601038
4. T. Banks, M. R. Douglas and N. Seiberg, “Probing F-Theory With Branes”, Phys. Lett. **387B** (1996) 278, hep-th/9605199
5. O. Aharony, J. Sonnenschein, S. Theisen and S. Yankielowicz “Field Theory Questions for String Theory Answers” hep-th/9611222
6. J. Polchinski, “Dirichlet Branes and Ramond-Ramond Charges”, Phys. Rev. Lett. **75** (1995) 4724, hep-th/9510169
7. J. Polchinski, S. Choudhuri and C. Johnson, hep-th/9502052
8. E. Witten, “Small Instantons in String Theory”, Nucl. Phys. **B460** (1996) 541, hep-th/9511030
9. E. Witten, “Bound States of Strings and p -branes”, Nucl. Phys. **B460** (1996) 335, hep-th/9510135
10. B. Greene, A. Shapere, C. Vafa and S. -T. Yau Nucl. Phys. **B337** (() 1990) 1.
11. D. R. Morrison and C. Vafa, “Compactifications of F-Theory on Calabi-Yau Three-folds I,II”, Nucl. Phys. **B473** (1996) 74, hep-th/9602114; Nucl. Phys. **B476** (1996) 437, hep-th/9603161
12. M. Bershadsky, K. Intriligator, S. Kachru, D. R. Morrison, V. Sadov and C. Vafa, “Geometric Singularities and Enhanced Gauge Symmetries”, hep-th/9605200
13. A. Sen “A Non-perturbative Description of the Gimon-Polchinski Orientifold” hep-th/9611186 Nucl. Phys. **B489** (() 1997) 139; “F-theory and the Gimon-Polchinski Orientifold” hep-th/9702061; “Orientifold Limit of F-theory Vacua” hep-th/9702165.
14. This topic was also discussed in: M. R. Douglas, D. A. Lowe and J. H. Schwarz “Probing F-theory With Multiple Branes” hep-th/9612062 Phys. Lett. **394B** (1997) 297
15. M. Bershadsky and A. Johansen, “Colliding Singularities in F Theory and Phase Transitions”, hep-th/9610111
16. J. D. Blum and A. Zaffaroni, “An Orientifold from F-Theory”, Phys. Lett. **387B** (1996) 71, hep-th/9607019
17. A. Dabholkar and J. Park, “A Note on Orientifolds and F-Theory”, hep-th/9607041
18. R. Gopakumar and S. Mukhi, “Orbifold and Orientifold Compactifications of F-Theory and M Theory to Six and Four Dimensions”, hep-th/9607057
19. R. G. Leigh and M. Strassler, “Exactly Marginal Operators and Duality in Four Dimensional $N = 1$ Supersymmetric Gauge Theory”, Nucl. Phys. **B447** (1995) 95, hep-th/9503121
20. O. Aharony, S. Kachru and E. Silverstein, “New $N = 1$ superconformal field theories in four dimensions from D-brane probes”, hep-th/9610205